

Original Article

Theophylline and silymarin are potent inhibitors of IL-4 secretion from activated human basophils

Hamidreza Jahanbani-Ardakani¹, Ghazale Bordbari², Nahid Eskandari^{2*}, Ali Mohammad Sabzghabae³, Alireza Andalib²

¹Isfahan Medical Students Research Center (IMSRC), Isfahan University of Medical Sciences, Isfahan, Iran

²Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

³Isfahan Clinical Toxicology Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract. Basophils are the most important cells in allergic reactions and secrete interleukin 4 (IL-4) and tumor necrosis factor α (TNF α) mediators. Theophylline as a phosphodiesterase 4 inhibitor and silymarin with its anti-inflammatory effects seem to be useful drugs in treatment of allergic reactions. The aim of the present study was comparing effects of theophylline and silymarin on IL-4 and TNF α secretion by human basophils. After isolation of peripheral blood mononuclear cells, three cultures were designed; theophylline, silymarin and control cultures. Cultures were incubated for 15 minutes with theophylline, silymarin and buffer before challenging with bacterial lipopolysaccharide and anti-IgE antibody. IL-4 and TNF α mediators were assessed by ELISA assay and values of each culture were compared with control group. Silymarin and theophylline as potent inhibitors decreased secretion of IL-4 from human basophils significantly ($P < 0.01$ and $P < 0.01$, respectively). However, neither theophylline nor silymarin had significant effect on TNF α secretion ($P = 0.14$ and $P = 0.08$, respectively). These data suggest that silymarin and theophylline could regulate the generation of IL-4 from human basophils.

Keywords: Theophylline, silymarin, interleukin-4, tumor necrosis factor α , basophil

Introduction

Basophils are well known for their pivotal role in allergic reactions. These cells are at least abundant leukocytes in peripheral blood (less than 1 percent). Basophils like mast cells have high-affinity IgE receptors on cell surface and also contain granules in their cytoplasmic space that will be exocytosed when cross-linked antigen-specific IgE occurred. Granules consist of histamine, leukotrienes, cytokines, chemokines and other mediators that are responsible for early and late allergic reactions [1]. In allergic disorders, basophils promote T helper2 (Th2) differentiation by IL-4 production and other co-stimulatory molecules such as CD40, CD80 and CD86 [2, 3]. Basophils would consider as a key role in allergic reaction for major source of cytokines such as IL-4 and Tissue necrosis factor alpha (TNF α) after activation through IgE receptors [4-6]. Production of these cytokines could be induced either by IgE-dependent mechanism or non-IgE dependent mechanism.

Silymarin is a flavonoid derived from milk thistle, *Silybum marianum* L. It is an antioxidant agent with protective effects on liver organ and prescribed clinically in liver diseases [7, 8]. In addition, silymarin has been

described to represent antioxidant, immunomodulatory, anti-proliferative, anti-fibrotic, and antiviral activities. The mechanisms of silymarin is elusive and its clinical efficacy is currently uncertain. Several studies, have reported immunomodulatory actions of silymarin *in vitro*. In addition, silymarin could modulate immune system and stabilize cell membrane. Previously, it was shown that silymarin could reduce histamine release from basophils and also promotes of Th2 cytokines and suppresses of Th1 cytokines [9, 10]. However, data on its effects on Th1/Th2 balance is in controversy.

Cyclic monophosphate phosphodiesterases (PDE) are an enzyme family that degrade cyclic adenosine monophosphate (cAMP) and guanosine monophosphate. PDE-4 enzyme, is a member of PDE enzyme that exist mostly in immune cells including neutrophils, monocytes, eosinophils and lymphocytes. Also, PDE-4 express highly in epithelial cells of lung and play a crucial role in pathogenesis of inflammatory lung diseases such as chronic obstructive pulmonary disease (COPD). PDE inhibitors prevent degradation of cyclic adenosine monophosphate (cAMP) and guanosine monophosphate. Moreover, PDE-4 inhibition results to accumulation of

* Corresponding author: Nahid Eskandari, MD, PhD
(neskandari@med.mui.ac.ir)




Figure 1 Effects of theophylline and silymarin on IL-4 secretion. Cells were incubated for 15 minutes before challenging with anti-IgE and LPS for 4 hours (n=5). Mean value of IL-4 was 60.06 ± 10.73 and 31.83 ± 12.43 in cultures treated with theophylline and silymarin respectively.

cAMP and consequently leads to suppressing expression of pre-inflammatory cytokines and chemokines [11, 12].

Theophylline (1, 3 dimethylxanthine) is a non-selective PDE inhibitor with anti-inflammatory and bronchodilator effects. Previously, it has been demonstrated that theophylline administration could lead to statistically improvement of forced expiratory volume in 1 second, forced vital capacity, and FEV1/FVC [13]. Also, long term administration of theophylline leads to reduction of interleukin 8 and tissue necrosis factor alpha in patients with chronic obstructive pulmonary disease [14]. Furthermore, theophylline administration in asymptomatic atopic asthmatic patients could reduce the serum level of IL-4 and also interleukin 5 [15].

The aim of the present study was to determine whether silymarin and theophylline regulates the generation of the cytokines, IL-4 and TNF α , from activated basophils.

Materials and Methods

Preparation of inhibitors and stimuli

Silymarin (Sigma, Poole, U.K.) was dissolved in 100% dimethyl sulfoxide (Sigma, Poole, U.K.) to achieve a 100mM stock solution. We used 10^{-2} concentration of silymarin stock solution. Theophylline (Sigma, Poole, UK) was prepared daily as stock solutions (2 mM) in the buffer. Escherichia coli bacterial lipopolysaccharide (LPS) (Enzo Life Sciences, Exeter, UK) was diluted with sterile HEPES buffer (2000ng/ml). Neat polyclonal goat anti-human IgE antibody (Sigma, Poole, UK) was made up in distilled water and stored at 4°C. All drugs were diluted to desired concentration in the buffer just before use.

Isolation of human basophils

Human peripheral blood mononuclear cells (PBMCs) were collected by vein puncture in heparinized tubes from whole blood obtained from fifteen healthy volunteers, who gave informed written consent. The study protocol was

approved by Ethics Committee of Isfahan University of Medical Sciences. Our exclusion criteria were as follows: any acute or chronic infectious disease, any neurological disorder, addiction and current smokers.

The cells were isolated by using standard Ficoll density-gradient centrifugation. Shortly, heparinized blood (50ml) was mixed with phosphate-buffered saline (PBS) (50ml) and after coating samples with Ficoll, the centrifugation (2800 rpm for 20 min) was performed and the interface layer of PBMCs was harvested and washed twice with PBS. Cell viability was assessed by trypan blue dye exclusion (0.4% trypan blue in PBS). Cells with viability of 95% and more were selected for further experiments.

Cell culture and mediator release

The cells were categorized in three cultures; control, theophylline and silymarin cultures. The cultures contained RPMI 1640 buffer (Gibco BRL, Dundee, U.K.) supplemented with BSA (Sigma, Poole, U.K.) (1 mg/ 1ml), gentamicin (Gibco BRL, Dundee, U.K.) (10 mg /ml) and calcium chloride (made up to 1 mM). Cultures were incubated with theophylline, silymarin and buffer for duration of 15 minutes before challenging with LPS and anti-IgE antibody. Cells incubated in buffer alone were considered as values of spontaneous mediator release, and all values cited for other cultures were corrected by subtracting values of spontaneous mediator release.

In experiments monitoring mediator generation, basophils were activated for 4 hours to assess IL-4 generation and 24 hours to assess TNF α generation. These conditions for optimal generation of IL-4 and TNF α generation have been reported by others [16].

After activation, the cells were centrifuged at 450 g for 4 min and the supernatants saved and analyzed for mediator release. The supernatant was assayed for IL-4 and TNF α mediators by available commercial enzyme linked immunosorbent assay (ELISA) kit (Biolog Life Science Institute, Bremen, Germany). The OD of samples was measured at 450 nm using a Dynatech plate reader.

Statistical analysis

Wilcoxon test were applied for comparing data between study groups. Data are expressed as means $\pm$ standard deviation. A P value of less than 0.05 was considered statistically significant. All of the analysis was performed by using SPSS software (SPSS Inc., Chicago, IL, USA).

Results

Effects of theophylline and silymarin on IL-4 secretion

The means $\pm$ SEM generation of IL-4 concentration inhibited by theophylline and silymarin were 5.68 ± 1.53 ng/mL and 9.69 ± 1.77 ng/mL respectively. Both theophylline and silymarin had significant effects on reduction of IL-4 secretion ($P < 0.01$ and $P < 0.01$ respectively) (Figs. 1 and 2). As figures 1 depicts, theophylline is more effective inhibitor than silymarin.




Figure 2 Effects of theophylline and silymarin on TNF α secretion. Cells were incubated for 15 minutes before challenging with anti-IgE and LPS for 24 h (n=5). Mean value of TNF α was 18.08 ± 2.61 and 22.93 ± 1.94 in cultures treated with theophylline and silymarin, respectively.

Effects of theophylline and silymarin on TNF α secretion

The mean value of TNF α inhibition in silymarin and theophylline cell cultures were 1.53 ± 0.04 ng/mL and 1.62 ± 0.52 ng/mL respectively. Neither silymarin nor theophylline had significant effects on reduction of TNF α secretion ($P=0.08$ and $P=0.14$ respectively) (Fig. 2).

Discussion

In the present study, we have investigated the effects of silymarin and theophylline on IL-4 and TNF α generation from human basophils.

A recent study showed topical administration of silymarin in NC/Nga mice with atopic dermatitis showed significant decrement of serum level of IL-4 [16]. Bakhshaei et al. have shown that prescription of silymarin with anti-histamine had significantly better improvement in clinical symptom severity of allergic disorders and the level of IL-4 decreased numerically in the sera of patients [17]. In the present study, it was shown that silymarin is effective inhibitor of the IL-4 generation from basophils, confirming observations made by others [16]. Previously inhibition of histamine releasing by theophylline from basophils has been reported [18]. Our data demonstrate the inhibitory effects of theophylline on IL-4, as reported by others [19, 20].

Further studies were performed to determine silymarin and theophylline effects that regulate cytokine generation from human basophils. Both compounds inhibited the IgE-mediated generation of TNF α . Earlier, it has been shown that long term treatment with theophylline in COPD patients lead to significant reduction of TNF α in sputum and could be associated with FEV1 increase [21]. In addition, theophylline could enhance basophil apoptosis and maybe this effect is one of the explanations of its affectivity in allergic diseases [9].

The mechanism of theophylline on decrement of pro-

inflammatory cytokines generation is inhibition of NF-kappaB activation via protection of IkappaBalpha protein [19]. Many properties of silymarin has been reported; some of them are anti-oxidant activity, anti-inflammatory activity, immuno-suppressive and immunomodulatory effects [22, 23]. The large body of data showed that Silymarin increases lymphocyte proliferation, interferon gamma (IFN- γ), IL-4 and IL-10 secretion by activated lymphocytes, in a dose-dependent manner [24]. Previously Han and colleagues reported that silymarin inhibits chemical-induced irritant contact dermatitis in mice [25].

In individual studies, administration of silymarin in patients with peritoneal dialysis leads to decreasing in serum level of TNF α . In another study in patients with β -thalassemia major it was shown that silymarin could significantly decrease TNF α in serum of patients [26, 27].

Inhibition of TNF α expression by silymarin had been reported by Han et al. and Zi and colleagues [25, 28]. The recent meta-analysis showed that administration of silymarin in patients with hepatitis B can decrease TNF α , TGF- β 1 and IL-6 [29]. Our results showed numerically but not significantly decrement of TNF α secretion by human basophils. This data could suggest that silymarin affects other sources of TNF α secretors except basophils; however more studies are suggested in this regard.

Silymarin exerts its anti-inflammatory effects via phosphorylation of IkBa and consequently it leads to inhibition of NF- κ B. NF- κ B influence production of cytokines, chemokines and other mediators related to inflammation [30, 31]. Therefore it could be concluded that PDE-4 inhibitors exerts their effect via inhibition on NF- κ B and due to wide range of activity of NF- κ B. It seems that more inflammatory mediators will be affected by PDE-4 inhibitors treatment.

Our study limitations were low number of cell cultures due to financial problems and lack of assessment of other involved cytokines in allergic reactions such as interleukin13 and interleukin 5. We recommend future studies to investigate inhibitory effects of silymarin and theophylline on IL-6, IL-13, IL-25 and other involved biomarkers in allergic reactions secreted from human basophils.

Conclusion

In conclusion, the present study demonstrates significant effects of silymarin on mediator releasing from human basophils, and it seems that theophylline and silymarin could be useful drugs in allergic diseases.

Acknowledgment

This project was supported by Isfahan University of Medical Sciences (Grant No: 393412). Also, the authors would like to greatly appreciate Dr. Abolfazl Fallah for his kind help.

Conflicts of interest

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

References

1. Siracusa MC, Kim BS, Spergel JM, Artis D. Basophils and allergic inflammation. *J Allergy Clin Immunol* 132:789-801, 2013.
2. Sokol CL, Medzhitov R. Role of basophils in the initiation of Th2 responses. *Cur Opin Immunol* 22:73-77, 2010.
3. Sokol CL, Chu N-Q, Yu S, Nish SA, Laufer TM, Medzhitov R. Basophils function as antigen-presenting cells for an allergen-induced T helper type 2 response. *Nature Immunol* 10:713-720, 2009.
4. Schneider E, Thieblemont N, De Moraes ML, Dy M. Basophils: new players in the cytokine network. *Eur Cytokine Network* 21:142-153, 2010.
5. Steiner M, Huber S, Harrer A, Himly M. The evolution of human basophil biology from neglect towards understanding of their immune functions. *BioMed Res Int* 8232830, 2016.
6. Gibbs BF. Basophils as key regulators of allergic inflammation and Th2-type immunity. *World Allerg Org J* 1:123-128, 2008.
7. Bannwart CF, Peracoli JC, Nakaira-Takahagi E, Peracoli MT. Inhibitory effect of silibinin on tumor necrosis factor- α and hydrogen peroxide production by human monocytes. *Natural Prod Res* 24:1747-1757, 2010.
8. Dixit N, Baboota S, Kohli K, Ahmad S, Ali J. Silymarin: A review of pharmacological aspects and bioavailability enhancement approaches. *Indian J Pharmacol* 39:172-179, 2007.
9. Kawakami A, Suzukawa M, Koketsu R, Komiya A, Ohta K, Yamamoto K, et al., editors. Enhancement of basophil apoptosis by olopatadine and theophylline. *Allergy and Asthma Proceedings*. OceanSide Publications, Inc., 2008.
10. Choi YH, Yan GH. Silibinin attenuates mast cell-mediated anaphylaxis-like reactions. *Biolog Pharmaceut Bulletin* 32:868-875, 2009.
11. Serezani CH, Ballinger MN, Aronoff DM, Peters-Golden M. Cyclic AMP: master regulator of innate immune cell function. *Am J Resp Cell Mol Biol* 39:127-132, 2008.
12. Page CP, Spina D. Selective PDE inhibitors as novel treatments for respiratory diseases. *Cur Opin Pharmacol* 12:275-286, 2012.
13. Margay SM, Farhat S, Kaur S, Teli HA. To study the efficacy and safety of doxophylline and theophylline in bronchial asthma. *J Clin Diag Res* 9:Fc05-08, 2015.
14. Iiboshi H, Ashitani J, Katoh S, Sano A, Matsumoto N, Mukae H, et al. Long-term treatment with theophylline reduces neutrophils, interleukin-8 and tumor necrosis factor- α in the sputum of patients with chronic obstructive pulmonary disease. *Pulm Pharmacol Ther* 20:46-51, 2007.
15. Kosmas E, Michaelides S, Polychronaki A, Roussou T, Toukmatzi S, Polychronopoulos V, et al. Theophylline induces a reduction in circulating interleukin-4 and interleukin-5 in atopic asthmatics. *Eur Resp J* 13:53-58, 1999.
16. Kang JS, Yoon WK, Han MH, Lee H, Lee CW, Lee KH, et al. Inhibition of atopic dermatitis by topical application of silymarin in NC/Nga mice. *Int Immunopharmacol* 8:1475-1480, 2008.
17. Bakhshae M, Jabbari F, Hoseini S, Farid R, Sadeghian MH, Rajati M, et al. Effect of silymarin in the treatment of allergic rhinitis. *Otolaryng Head Neck Surg* 145:904-909, 2011.
18. Nakano J, Yano T, Yamamura K, Yoshihara H, Ohbayashi O, Yamashita N, et al. Aminophylline suppress the release of chemical mediators in treatment of acute asthma. *Respir Med* 100:542-550, 2006.
19. Chen ZC, Ding JJ, Qian HY, Lin XH, Feng KQ, Wang J. Clinical efficacies of inhaled corticosteroids plus theophylline in the treatment of bronchial asthma]. *Zhonghua Yi Xue Za Zhi* 93:1076-1079, 2013.
20. Eskandari N, Wickramasinghe T, Peachell PT. Effects of phosphodiesterase inhibitors on interleukin-4 and interleukin-13 generation from human basophils. *Brit J Pharmacol* 142:1265-1272, 2004.
21. Iiboshi H, Ashitani J-i, Katoh S, Sano A, Matsumoto N, Mukae H, et al. Long-term treatment with theophylline reduces neutrophils, interleukin-8 and tumor necrosis factor- α in the sputum of patients with chronic obstructive pulmonary disease. *Pulm Pharm Therap* 20:46-51, 2007.
22. Min K, Yoon W-K, Kim SK, Kim B-H. Immunosuppressive effect of silibinin in experimental autoimmune encephalomyelitis. *Arch Pharm Res* 30:1265-1272, 2007.
23. Kang JS, Park S-K, Yang K-H, Kim HM. Silymarin inhibits TNF- α -induced expression of adhesion molecules in human umbilical vein endothelial cells. *FEBS Letters* 550:89-93, 2003.
24. Alidoost F, Gharagozloo M, Bagherpour B, Jafarian A, Sajjadi SE, Hourfar H, et al. Effects of silymarin on the proliferation and glutathione levels of peripheral blood mononuclear cells from beta-thalassemia major patients. *Int Immunopharmacol* 6:1305-1310, 2006.
25. Han MH, Yoon WK, Lee H, Han S-B, Lee K, Park S-K, et al. Topical application of silymarin reduces chemical-induced irritant contact dermatitis in BALB/c mice. *Int Immunopharmacol* 7:1651-1658, 2007.
26. Gharagozloo M, Karimi M, Amirghofran Z. Immunomodulatory effects of silymarin in patients with β -thalassemia major. *Int Immunopharmacol* 16:243-247, 2013.
27. Nazemian F, Karimi G, Moatamedi M, Charkazi S, Shamsara J, Mohammadpour AH. Effect of silymarin administration on TNF- α serum concentration in peritoneal dialysis patients. *Phytother Res* 24:1654-1657, 2010.
28. Esmaeil N, Anaraki SB, Gharagozloo M, Moayed B. Silymarin impacts on immune system as an immunomodulator: One key for many locks. *Int Immunopharmacol* 50:194-201, 2017.
29. Wei Q, Xu X, Ling Q, Zhou B, Zheng S-S. Perioperative antiviral therapy for chronic hepatitis B-related hepatocellular carcinoma. *Hepatobil Pancreat Dis Int* 12:251-255, 2013.
30. Kim B-R, Seo H-S, Ku J-M, Kim G-J, Jeon CY, Park JH, et al. Silibinin inhibits the production of pro-

inflammatory cytokines through inhibition of NF- κ B signaling pathway in HMC-1 human mast cells. *Inflamm Res* 62:941-950, 2013.

31. Striz I, Brabcova E, Kolesar L, Liu X, Barcova

I, Sekerkova A, et al. Epithelial cells modulate genes associated with NF kappa B activation in co-cultured human macrophages. *Immunobiology* 216:1110-1116, 2011.